

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963 A

Report SAM-TR-82-5



RESPONSES TO HARMONIC ACCELERATION WITH VARYING HEAD POSTURES

Daniel E. Dreher, M.A. Kennith W. Stevens, B.S. Edward J. Engelken, M.S. James W. Wolfe, Ph.D.

SELECTED

March 1982

Final Report for Period 1 March 1981 - 30 October 1981

Approved for public release; distribution unlimited.

USAF SCHOOL OF ARROSPACE MEDICINE Acrospace Medical Division (AFSC) Breeks Air Force Base, Texas 78255

08 24



This final report was submitted by personnel of the Vestibular Function Otolaryngology Branch, Clinical Sciences Division, and the Data Sciences Mivision, USAF School of Aerospace Medicine, Aerospace Medical Division, AFSC, Brooks Air Force Base, Texas, under job order 7755-17-41.

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation the Government thereby incurs no responsibility nor any obligation whatsdown; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be repair by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The voluntary informed consent of the subjects used in this research was obtained in accordance with AFR 169-3.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

Tanul 4. DANIEL E. DREHER, M.A.

Project Scientist

amie Withake JAMES W. WOLFE, Ph.D. Supervisor

Colonel, USAF, MC Commander

Accession For NTIS PRA&I DTIC TAB Unannounced Justification Distribution/ Availability Codes Avail and/or Special

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
SAM-TR-82-5	AD-AII4 %	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
RESPONSES TO HARMONIC ACCELERATION	WITH	Final Report
VARYING HEAD POSITIONS		1 Mar 1981 - 30 Oct 1981
		6. PERFORMING ORG, REPORT NUMBER
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Daniel E. Dreher, M.A.; Kennith W.	Stevens, B.S.:	
Edward J. Engelken, M.S.; James W.		
9 PERFORMING ORGANIZATION NAME AND ADDRESS USAF School of Aerospace Medicine (NC F DD)	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Aerospace Medical Division (AFSC)	NG Q DK)	62202F
Brooks Air Force Base, Texas 78235		7755-17-41
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
USAF School of Aerospace Medicine (NG & BR)	March 1982
Aerospace Medical Division (AFSC)		13. NUMBER OF PAGES
Brooks Air Force Base, Texas 78235		5
14. MONITORING AGENCY NAME & ADDRESS(It different	from Controlling Office)	15. SECURITY CLASS. (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, it different from Report) 18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Harmonic acceleration Head position		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
To determine if a particular head position results in maximal response to stimulation, 50 subjects were tested with sinusoidal acceleration in three head positions with reference to the horizontal plane. A PDP-11/34 digital computer was used to analyze the responses. The pertinent parameters were gain, phase, and directional preponderance. The results showed no difference in the parameters for the three head positions. This suggests that positioning a subject's head for rotational testing is not as critical as previously believed.		
DD FORM 1473 FRITION OF L NOV 65 IS ORSOLETE		

DD $_{1}$ $_{1}$ $_{1}$ $_{1}$ $_{2}$ $_{3}$ $_{1}$ $_{2}$ $_{3}$ $_{4}$ $_{2}$ $_{3}$ $_{4}$ $_{2}$ $_{3}$ $_{4}$ $_{5}$

UNCLASSIFIED

RESPONSES TO HARMONIC ACCELERATION WITH VARYING HEAD POSITIONS

INTRODUCTION

The major function of the vestibular system is to provide a stable platform for vision by initiating compensatory eye movements in response to head movements. The reflexive, compensatory eye movements—the vestibulo—oculomotor reflex (VOR)—function to keep an object in the visual field whenever the head is rotated, elevated, or tilted.

Classically, vestibular function has been tested using caloric stimulation. Within the last decade, however, advances in torque motor drive systems and minicomputers have led to increased use of rotational stimulation in clinical evaluations. We have used sinusoidal acceleration in this laboratory for several years to study the effects of peripheral and central pathology on the VOR (1-5) and the effects of surgical procedures used in treatment of vestibular disorders (6, 7) and to compare results of caloric and rotational testing (3, 8). In these studies, we assumed that maximum responses to rotational stimulation would occur when the subject's head was positioned with the chin down approximately 25°, the position where the canals are approximately in the plane of rotation. To the best of our knowledge, however, there is no experimental evidence which proves that this position produces maximum responses. This study was designed to determine if any head position consistently produces maximal responses to horizontal sinusoidal stimulation.

In 1938 Travis (9) studied the effects of various head positions on the subjective sensation of rotation. He used sinusoidal rotation and had his subjects indicate the intensity of their perception of motion when their heads were in three different positions: 0° , $+45^{\circ}$, and -45° , with respect to the plane of the horizontal canals. His subjects reported greater sensation when their heads were in the $+45^{\circ}$ position. He postulated that this occurred because the vertical canals were being stimulated along with the horizontal canals in the 45° position. We chose head positions of 0° and $\pm 15^{\circ}$ on the assumption that if a subject's head is improperly positioned for testing, the error is probably within just a few degrees of the 0° position. Angles of $\pm 15^{\circ}$ provide sufficient range to include any error that would probably occur in clinical testing.

METHODS

From on-base volunteers, we selected 50 subjects who reported a history free of vestibular problems. Each subject was instrumented with silver silver-chloride surface electrodes in the usual manner for recording eye movements and was seated in a chair mounted on a torque motor (Contraves-Goerz, Pittsburgh, Pa., Model DP-300) inside a lightproof room. Ten minutes were allowed for dark adaptation. The test procedure consisted of three trials of .04-Hz harmonic stimulation with a 5-minute break between each trial. A technician established the desired head position before each trial and adjusted a bite bar and headrest

so that the subject could maintain the position during rotation. Subject alertness was maintained by a tone-discrimination task. A Digital Equipment Company PDP-11/34 minicomputer provided the .04-Hz driving signal and analyzed the resulting eye-movement recordings. Each trial lasted approximately 4 minutes, with a peak velocity of $60^{\circ}/s$. The technician repositioned the subject's head for the next trial during the 5-minute breaks.

A program implemented on the PDP-11/34 computer digitized the eye-movement data and calculated eye relocity and eye acceleration (10). The fast-phase components of the tracings were identified using velocity and acceleration thresholds. Once identified, the fast-phase components were removed from the eye-velocity record by linear interpolation, resulting in a record containing only slow-phase eye velocity. Frequency domain analysis was performed on the stimulus signal and the slow-phase eye-response record, producing estimates of VOR gain, phase, and directional preponderance for each trial.

Gain is defined as the ratio of the magnitude of the slow-phase eye-velocity response to the magnitude of the head-velocity stimulus. Normally, gain is in the 0.4 to 0.6 range at the frequency used in this study. Phase describes the time relationship between the stimulus and the response, and directional preponderance indicates the left-right symmetry of the response. One study (11) indicates that gain is modulated by cortical function; however, by controlling subject alertness, we minimized all but the head velocity as factors affecting gain. Phase and directional preponderance are largely independent of VOR gain.

In this study the stimulus intensity (angular velocity of the head) was held constant, but the position of the horizontal canals with respect to the plane of rotation was varied. In this manner the canals received a varying degree of stimulation. The decrement in stimulus intensity received by the canals in the $\pm 15^{\circ}$ positions can be expressed in terms of the cosine function of the angle formed by the plane of rotation and the plane of the canals. Since the cosine of 15° is 0.96, we anticipated a 4% decrease in gain when the head was in the $\pm 15^{\circ}$ position.

RESULTS

The subjects had a mean age of 32 years, ranging from 19-55 years. Of the 50 subjects, 44 (88%) were males, 27 (54%) were nonflying military, 19 (38%) were aircrew members, 4 (8%) were civilians.

Analysis of variance showed no significant differences in gain, phase, or directional preponderance for the three head positions (p > .05). This was true even though phase lag and preponderance did increase in the $+15^{\circ}$ position. The variability of each parameter (gain, phase, and preponderance) was similar for the three head positions. Figure 1 shows these data. Our expectation that gain would be 4% higher in the 0° position was not supported by the following: 15 subjects (30%) had higher gain values in the 0° position, 16 (32%) had higher gain values in the -15° position, and 19 (38%) had higher gain values in the $+15^{\circ}$ position.




Figure 1. Head positions tested, with results for gain, phase, and preponderance.

DISCUSSION

Data from this study show that head position for rotational testing is not as critical as previously believed. Although technicians must attempt to align the horizontal canals with the plane of rotation, there is some allowance for a small margin of error and individual physiological differences. Recent anatomical studies (12, 13) have shown that the three canals on each side of the head are not exactly at right angles to each other. Also, the data from these studies confirm that the horizontal canal is tilted upward 25° (instead of the commonly accepted 30°) and that the medial side of the canal is higher than the lateral side. Often the planes of the two horizontal canals will differ by several degrees. Thus, positioning the head for rotational testing is a compromise: while one horizontal canal is in the plane of rotation, the other may be as much as 9° out of the plane (12). Because of this, responses to rotation must be accepted as a pattern of multicanal stimulation (13). We believe that our data reflect the influence of these individual anatomical variations.

ACKNOWLEDGMENTS

The authors wish to thank the following for their technical assistance: SSgt Kevin L. Dever, John W. Docken, Sgt Patricia A. Hladsky, and TSgt Lloyd J. Loup. We also wish to thank Dr. H. H. Hanna for editorial assistance and Rosa Linda Rodriguez for manuscript preparation.

REFERENCES

- 1. Wolfe, J. W., and C. M. Kos. Nystagmic responses of the rhesus monkey to rotational stimulation following unilateral labyrinthectomy. Trans Am Acad Ophthalmol Otolaryngol 84:38-45 (1977).
- 2. Wolfe, J. W., et al. Low-frequency harmonic acceleration as a test of labyrinthine function: Basic methods and illustrative cases. Trans Am Acad Ophthalmol Otolaryngol 86:130-142 (1978).
- 3. Wolfe, J. W., et al. Vestibular responses to bithermal caloric and harmonic acceleration in patients with Meniere's disease. Ann Otol Rhinol Laryngol 87(6):861-867 (1978).
- 4. Wolfe, J. W., et al. Low-frequency harmonic acceleration in the evaluation of patients with peripheral labyrinthine disorders. In Honrubia and Brazier (eds.). Nystagmus and Vertigo: clinical approaches to the patient with dizziness. New York: Academic Press, Inc. (In press).
- 5. Olson, J. E., and J. W. Wolfe. Comparison of subjective symptomatology and responses to harmonic acceleration in patients with Meniere's disease. Ann Otol Rhinol Laryngol 90:15-17, Suppl 86 (1981).
- 6. Wolfe, J. W., et al. Low-frequency harmonic acceleration in the evaluation of surgical treatment of Meniere's disease. Adv Otorhinolaryngol 25:192-196 (1979).

- 7. Olson, J. E., and J. W. Wolfe. Responses to low-frequency harmonic acceleration in patients with acoustic neuromas. Laryngoscope 91:1270-1277 (1981).
- 8. Wolfe, J. W., et al. Vestibular responses to bithermal caloric and harmonic acceleration in patients with Meniere's disease. In K. H. Vosteen et al. (eds.). Meniere's disease: New York: Theim Stratton, Inc., 1981.
- 9. Travis, R. C. The effects of varying the position of the head on voluntary response to vestibular stimulation. J Exp Psychol 23:295-303 (1938).
- 10. Engelken, E. J., et al. Application of digital filters in the processing of eye movement data. Behav Res Meth Instrum (In press).
- 11. Barr, C. C., et al. Voluntary, non-visual control of the human vestibuloocular reflex. Acta Otolaryngol 81:365-375 (1976).
- 12. Blanks, R. H., et al. Planar relationships of the semicircular canals in man. Acta Otolaryngol 80:185-195 (1975).
- 13. Curthoys, I. S., et al. Semicircular canal functional anatomy in cat, guinea pig, and man. Acta Otolaryngol 83:258-265 (1977).

